


Gluon and Ghost Propagators in Landau Gauge on the Lattice

A.Y. Lokhov^a, C. Roiesnel

Centre de Physique Théorique^b de l'Ecole Polytechnique, 91128 Palaiseau Cedex, France.



We study the ultraviolet behaviour of the ghost and gluon propagators in quenched QCD using lattice simulations. Extrapolation of the lattice data towards the continuum allows to use perturbation theory to extract Λ_{QCD} - the fundamental parameter of the pure gauge theory. The values obtained from the ghost and gluon propagators are coherent. The result for pure gauge SU(3) at three loops is $\Lambda_{\overline{\text{MS}}} \approx 320 \text{ MeV}$. However this value does depend strongly upon the order of perturbation theory and upon the renormalisation description of the continuum propagators. Moreover, this value has been obtained without taking into account possible power corrections to the short-distance behaviour of correlation functions.

1 Introduction

Correlation functions involving ghosts appear in Schwinger-Dyson equations in covariant gauge. Thus, these correlation functions play an important role in the non-perturbative studies of the infrared behaviour of the gluon propagator and running coupling constant, and of the confinement. Lattice simulations provide a non-perturbative way to obtain propagators and vertices in Landau gauge. The ghost propagator has not been studied¹ so much as the gluon propagator in the case of QCD ($SU(3)$ gauge group). That is why one should first probe the lattice formulation of the ghost propagator by studying its behavior at small distances, where lattice results may be checked using perturbation theory. Moreover, the perturbative fit of the ghost and gluon propagators gives us the value of Λ_{QCD} - the fundamental parameter of the pure gauge theory, which is a renormalisation group invariant, but is scheme-dependent. Even if this parameter is not a physical quantity, it can serve as a definition of the unique energy scale of quenched QCD. In this contribution we report on our on-going study of the ghost and gluon propagators in the domain of energy $2 \text{ GeV} \leftrightarrow 6 \text{ GeV}$. We have fitted these correlation functions

^aTalk given by Alexey Lokhov.

^bUnité Mixte de Recherche C7644 du CNRS

using three-loop perturbative expansion in the $\widetilde{\text{MOM}}$ renormalisation scheme⁷. This allows us to obtain two values for Λ_{QCD} , and thus to check the self-consistency of the lattice approach. In what follows we will describe briefly the lattice formulation in the case of quenched QCD and present preliminary results for two-points correlation functions.

2 Simulation Setup

On the lattice one works with an Euclidean formulation of QCD in discretised four-dimensional space (with lattice spacing a). In practical simulations one considers this theory in a finite volume ($V = L^4$), with periodic boundary conditions for the gluon field. Then one can evaluate numerically the functional integrals which define Green functions, using Monte-Carlo techniques. In our simulations we have used the standard Wilson action to produce gauge field configurations (more details may be found in reviews²). Then the gauge has to be fixed. Presently, the Landau gauge is the only covariant gauge which can be implemented efficiently on the lattice. The Landau gauge is fixed by finding a minimum of the functional

$$F_U(\{g(x)\}) = -\text{ReTr} \sum_{x,\mu} g(x) U_\mu(x) g^\dagger(x + e_\mu), \quad g(x) \in SU(3), \quad (1)$$

where the link variable is related to the gauge field $A_\mu^a(x)$ by $U_\mu(x) = \exp i A_\mu^a(x) t^a$, t^a are the generators of the gauge group. A local minimum defines explicitly the gauge transformation that transforms $\{U\}$ to $\{U^{(g)}\}$ so, that $U_\mu^{(g)}(x)$ satisfies the discretised Landau condition

$$\frac{\delta}{\delta g(x)} F_U(\{g(x)\}) = 0 \quad \Rightarrow \quad \text{ReTr} \sum_\mu \left[U_\mu^{(g)}(x) + U_\mu^{(g)\dagger}(x - e_\mu) \right] = 0. \quad (2)$$

Once the gauge is fixed, one can calculate numerically the gluon two-point correlation function

$$\langle A_\mu^a(x) A_\mu^b(x) \rangle = \frac{1}{Z} \int [\mathcal{D}U] A_\mu^a(x) A_\mu^b(y) e^{-S_{\text{Wilson}}[U]}, \quad (3)$$

where $A_\mu(x) = \frac{U_\mu(x) - U_\mu^\dagger(x)}{2i}$. In momentum space, and in the continuum limit, the general parametrisation of the gluon propagator in Landau gauge reads

$$\langle \tilde{A}_\mu^a(p) \tilde{A}_\nu^b(-p) \rangle = \frac{G(p^2)}{p^2} \delta^{ab} (\delta_{\mu\nu} - \frac{p_\mu p_\nu}{p^2}). \quad (4)$$

Thus, lattice simulations can provide non-perturbative information about the scalar function $G(p^2)$.

The ghost propagator in covariant gauge is defined in the continuum formulation of the QCD as the inverse of Faddeev-Popov operator

$$M^{ab}(x, y) = [\partial \cdot D]^{ab}(x, y). \quad (5)$$

where D is the covariant derivative in the adjoint representation. On the lattice this definition should be modified. Indeed, the Faddeev-Popov operator on the lattice is defined by the second derivative of the functional F_U at a local minimum

$$\frac{\delta^2}{\delta g(x) \delta g(y)} F_U(\{g(x)\}) \longrightarrow M_{\text{lattice}}^{ab}(x, y), \quad (6)$$

where (f^{abc} are the structure constants of the gauge group)

$$\begin{aligned} M_{\text{lattice}}^{ab}(x, y) = & \sum_\mu \left\{ G_\mu^{ab}(x) (\delta_{y, x+\hat{\mu}} - \delta_{y, x}) + G_\mu^{ab}(x - \hat{\mu}) (\delta_{y, x-\hat{\mu}} - \delta_{y, x}) \right. \\ & \left. + \frac{1}{2} \sum_c f^{abc} (\delta_{y, x+\hat{\mu}} A_\mu^c(x) - \delta_{y, x-\hat{\mu}} A_\mu^c(x - \hat{\mu})) \right\} \end{aligned} \quad (7)$$




Figure 1: Three-loop fit for the data obtained for $V = 32^2, \beta = 6.4$. The fit has been done in the interval $[2, 5\text{GeV}; 4\text{GeV}]$ for the gluon propagator (left image), and $[2, 5\text{GeV}; 4\text{GeV}]$ for the ghost propagator (right image).

and

$$G_\mu^{ab}(x) = \frac{1}{2} \text{Tr} \left(\left\{ t^a, t^b \right\} \left(U_\mu(x) + U_\mu^\dagger(x) \right) \right) \quad (8)$$

Inverting this operator with a local source at the origin with no zero-mode component

$$\left(1 - \frac{1}{V}, -\frac{1}{V}, \dots, -\frac{1}{V} \right)$$

(V is the volume of the lattice), and performing a Fourier transformation, one can extract the scalar form factor $F(p^2)$ of the ghost propagator in Landau gauge for all momenta at once

$$\langle c^a(p) \bar{c}^b(-p) \rangle = \frac{F(p^2)}{p^2} \delta^{ab}. \quad (9)$$

Let us discuss briefly the errors of the method. Obviously, one has the statistical error due to the Monte-Carlo calculation of the functional integral. The systematic error is mostly due to the space-time discretisation. Indeed, the rotational symmetry is broken down to the group H_4 - the symmetry group of the hypercube. That means that, at most momenta, the calculated values of any scalar quantity will split, because an orbit of the H_4 group is not uniquely defined by the value of p^2 . Thus, one should extrapolate all data points to the continuum limit $a \rightarrow 0$. Now we have a well developed method of performing such an extrapolation^{3,4}. All plots, presented further, have been done using this method. Another source for systematic error comes from the Euclidean formulation of the QCD and gauge fixing (lattice version of Gribov copies⁵). But the so-called Gribov noise is negligible in the short-distance study that is discussed here.

3 Results

We have done simulations on lattices with size $\{16^4, 24^4, 32^4\}$ (1000, 500, 250 configurations, respectively) and for the respective values of the lattice bare couplings $\beta = \{6.0, 6.2, 6.4\}$ ($\beta = \frac{6}{g_0^2}$, where g_0 is the value of the bare coupling at the cut-off energy scale a^{-1}). Thus, these lattices have roughly the same physical volume. We did a fit of both propagators at three loops in the $\overline{\text{MOM}}$ scheme⁷, which was shown⁴ to provide a good description of the lattice gluon propagator,

and then converted the value of $\Lambda_{\widetilde{MOM}}$ to the more familiar $\overline{MS}$ scheme. The results for $\Lambda_{\overline{MS}}$ are given in Table 1. Two errors are quoted. The error between parentheses is the statistical one and is rather small. The other error is the systematic error which can be quite large, especially for the ghost propagator where it can reach 10%. The systematic error has been estimated from the stability of the fits in the region [2GeV;6GeV]. Examples of the fits are given in Fig. 1.

Table 1: Results of the fits for Λ_{QCD}

β	L	$\Lambda_{\overline{MS}}^{(3)gluon}$	$\chi^2/\text{d.o.f}$	$\Lambda_{\overline{MS}}^{(3)ghost}$	$\chi^2/\text{d.o.f}$
6.0	16	$324(2)^{+2}_{-5}$ MeV	1.2	$322(8)^{+20}_{-16}$ MeV	0.5
6.2	24	$320(2)^{+8}_{-14}$ MeV	0.9	$326(5)^{+26}_{-33}$ MeV	0.6
6.4	32	$312(1)^{+9}_{-25}$ MeV	1.4	$331(4)^{+42}_{-35}$ MeV	0.9

4 Conclusion

We obtain coherent results for Λ_{QCD} at three loops from the ghost and gluon propagators. This means that the lattice QCD approach is self-consistent in the ultraviolet domain when studying correlation functions which involve ghosts. The value for Λ_{QCD} that we report here is approximatively equal to 320MeV in the $\overline{MS}$ scheme. However it should be emphasized that this value does depend strongly upon the order of perturbation theory used in the fits. This dependence manifests itself in two ways. First, two-loop perturbation theory is also able to reproduce the data but with a much higher Λ_{QCD} scale. Secondly, at any given order, the value of Λ_{QCD} relies very much on the renormalization scheme used to describe the propagators in the continuum. The value of the $\Lambda_{\overline{MS}}$ scale that we quote here should therefore be considered as an *effective* Λ_{QCD} scale.

In order to check the asymptoticity and the coherence of Λ_{QCD} more precisely one should also study the ultraviolet behaviour of the gluon and ghost propagators at four loops in several $\widetilde{MOM}$ schemes. Possible power corrections⁶ could also make the value of the Λ_{QCD} scale lower. We currently work on this, and the results will be reported elsewhere.

Acknowledgements

We thank our colleagues of the LPT-Orsay, P. Boucaud, J.P. Leroy, A. Le Yaouanc, J. Micheli and O. Pène, for many useful discussions.

References

1. H. Suman and K. Schilling, Phys. Lett. B **373** (1996) 314 [arXiv:hep-lat/9512003].
2. A. Sternbeck, E. M. Ilgenfritz, M. Muller-Preussker and A. Schiller, arXiv:hep-lat/0412011.
3. I. Montvay, G. Munster, “Quantum Fields on a Lattice,” Cambridge Univ. Press (1994).
4. D. Becirevic, P. Boucaud, J. P. Leroy, J. Micheli, O. Pène, J. Rodriguez-Quintero and C. Roiesnel, Phys. Rev. D **60** (1999) 094509 [arXiv:hep-ph/9903364].
5. D. Becirevic, P. Boucaud, J. P. Leroy, J. Micheli, O. Pène, J. Rodriguez-Quintero and C. Roiesnel, Phys. Rev. D **61** (2000) 114508 [arXiv:hep-ph/9910204].
6. T. D. Bakeev, E. M. Ilgenfritz, V. K. Mitrjushkin and M. Mueller-Preussker, Phys. Rev. D **69** (2004) 074507 [arXiv:hep-lat/0311041].
7. P. Boucaud *et al.*, JHEP **0004** (2000) 006 [arXiv:hep-ph/0003020].
8. K. G. Chetyrkin and A. Retey, “Three-loop three-linear vertices and four-loop MOM beta functions in massless QCD,” arXiv:hep-ph/0007088.